



**PROPOSAL FOR ECOLOGICAL CORRIDORS AND BUFFER ZONES AS MEASURES TO RESTORE THE CONNECTIVITY OF THE HABITAT OF THE SPECIES '*santamartamys rufodorsalis*' IN COLOMBIA**

**Proposal for Ecological Corridors and Buffer Areas as Measures to Restore the Connection in the Habitat of the '*santamartamys rufodorsalis*' Species in Colombia**

**How To Cite This Article:** COLOMBIA WILD CORPORATION, 2021. Proposal For Ecological Corridors and Buffer Zones as Measures To Restore The Connectivity Of The Habitat Of The Species '*Santamartamys Rufodorsalis*' In Colombia. COLOMBIA WILD CORPORATION. Santa Marta, Colombia. 18 pag.

**SUMMARY**

In the present study, the design of ecological corridors and buffer zones is proposed as a measure for the expansion and restoration of the connectivity of the habitat of the game species *Santamartamys rufodorsalis* in the north center of the Sierra Nevada of Santa Marta, located in north of Colombia. The methodology used, the result of modifications of the proposal by Bentrup (2008), was based on the analysis of geographic parameters that define areas with the best territorial suitability for the habitat of the species. Likewise, the study is based on a cost analysis that, through a Geographic Information System, allows defining the potential location of the functional connectors. In this way, the model provides reliable information that allows the exact layout of corridors and buffer zones to be proposed. As will be seen in due course, the methodology used can be adjusted for other species.

**Keywords:** ecological corridors; buffer zones; *Santamartamys rufodorsalis*; Colombia.





## **INTRODUCTION**

The fragmentation of ecosystems and the loss of ecological connectivity, two of the main factors that have a profound impact on biodiversity, are largely caused by the increase and homogenization of agricultural and urban activities (Rodríguez-Soto et al., 2013; De León Mata et al., 2014; Johnstone et al., 2014; Gao Q, 2014; García-Marmolejo et al., 2015; Loro et al., 2015; Van Langevelde, 2015; Villemey et al., 2015 ; Burkart et al., 2016; Peled, 2016). Ecological connectivity is defined according to the degree to which the territory facilitates or hinders significant ecological processes such as the movement of species through existing habitat resources in the landscape (San Vicente, 2014). Therefore, the preservation of ecological connectivity helps to minimize the negative effects of habitat fragmentation (Johnstone et al., 2014).

One way to contribute to reducing the loss of ecological connectivity consists in the design of ecological corridors that function as buffer zones against the negative effects of fragmentation. In this sense, it is vital to consider the theoretical insights of Bentrup (2008), who defines ecological corridors as “vegetation strips incorporated into the landscape [that serve] to influence ecological processes and provide a variety of goods and services. They are known by various names, such as wildlife corridors, greenways, windbreaks and filter strips” (p.1). However, other authors such as Tres and Reis (2007) maintain that many of these models have a strong tendency to fill the areas with tree species that limit the space available for natural regeneration. However, it should be added that the success of ecological corridors depends on the degree to which their design contributes to eliminating the biotic and abiotic barriers that degrade the ecosystem.

Although Colombia is characterized by having a high biodiversity, according to the red list of endangered animals it is one of the countries with the highest number of threatened species in the world (International Union for Conservation of Nature, 2014). Among the factors that accentuate this situation, it is worth highlighting the intense deforestation of forests and the overexploitation of natural resources due to hunting and illegal animal trafficking (Sierra, 2013; Rodríguez and Ortega, 2012; De Osma Vargas-Machuca et al., 2014). These circumstances contribute in various ways to the reduction and fragmentation of the habitat of game species such as the *Cuniculus paca*, known locally as Red Crested Tree Rat .

The Red Crested Tree Rat has a wide geographical distribution in the neotropical region of the Colombian territory: it inhabits the humid, dry, tropical and subtropical forests of the Coast and the Amazon, as well as some Andean foothills below 700 meters above sea level (Albuja et al. , 1993; Tirira, 2007). According to the Convention on International Trade in Endangered Species of Wild Fauna and Flora (2009), the species is not in danger of extinction; however, its survival in northwestern Colombia is threatened by multiple factors, among which its own reproductive characteristics stand out; the loss and fragmentation of the native forest — its natural habitat— due to the effects of deforestation, and the increase in poaching, given that it is used as food by the indigenous and rural populations of the region (Tirira, 2007, 2011; Rodríguez and Ortega, 2012; Bonilla-Morales et al., 2013; Pozo, 2013; De Osma Vargas-Machuca et al., 2014).

As stated in the study by De Osma Vargas-Machuca et al. (2014), despite their decline and fragmentation, the forests continue to be the habitat of the Red Crested Tree Rat or *Cuniculus paca*, and to a lesser extent other game species such as *Dasyprocta punctata* or *guatusa* and the *Dasyopus novemcinctus* or armadillo. These species are located in the remnants of native forests located in the northwest of Manabí.

Accordingly, the objective of this research is to propose the design of ecological corridors and buffer zones in the sectors with the best territorial suitability for Red Crested Tree Rat ; surfaces defined according to socio-economic, biophysical and social factors. Thus, the purpose of the project is to provide adequate measures to restore ecological connectivity between the remnants of native forest and thus favor an increase in the habitat of the species. In particular, it seeks to facilitate the mobility of the Red Crested Tree Rat in a sector that covers 1,500 km<sup>2</sup> in the center-north of the province of Manabí: a space for agricultural, silvopastoral, and agroforestry activities.




**Proposal for ecological corridors and buffer zones as measures to restore the connectivity of the habitat of the species *Santamartamys rufodorsalis* in Colombia**

CC-052-2019

Version 3

June 01, 2020

**PROJECT MAP**




**ECOLOGICAL CONNECTIVITY OF THE RED CRESTED TREE RAT (*Santamartamys rufodorsalis*)**

	Investigation area
	Priority habitats of target species
	Area with high poaching threat
	Arahuc indigenous cities

**Scale**

1cm=50,000 meters



**Figure 1. Location of the study area**

Cartographic source: base map of the Military Geographic Institute (2010) and thematic maps of the Ministry of the Environment of Colombia (2014)



**Proposal for ecological corridors and buffer zones as measures to restore the connectivity of the habitat of the species *Santamartamys rufodorsalis* in Colombia**

**CC-052-2019**

**Version 3**

**June 01, 2020**

Although it is not under a specific political-administrative delimitation, the study area includes part of the cantons of Flavio Alfaro, Chone and El Carmen. The surface is located between the following coordinates: 00° 17' 03" S and 80° 01' 24" W as a northwestern point; 00° 17' 03" S and 79° 38' 31" W as northeast point; 00° 26' 45" S and 80° 01' 24" W as the southwest point, and 00° 26' 45" S and 79° 38' 31" W as the southeast point (figure 1).

## **METHODOLOGY**

For the formulation of a proposal for ecological connectors in the study area, it was necessary to a) identify the native forest cover; b) identify the territorial aptitude for the presence of Red Crested Tree Rat (*Cuniculus paca*); c) propose the creation of connectors —ecological corridors and buffer zones—, and d) process geoinformation (figure 2).

### **Identification of native forest cover**

Through the visual analysis provided by the ArcGIS Geographic Information Systems package (version 10.1), the forest cover was interpreted and delimited. The procedure was developed on a set of orthophoto maps at a scale of 1:5000 generated by the Ministry of Agriculture through the Program of the National System of Information and Management of Rural Lands and Technological Infrastructure (Sigtierras), under the supervision of the Military Geographic Institute (IGM). The orthophoto maps correspond to a UTM coordinate system, WGS-84 datum, Zone 17. Likewise, these coordinates were acquired in 2015.


### **Identification of variables and preparation of maps of optimal areas by criteria**

To develop the research, firstly, factors, variables and criteria were defined in a matrix that represented the cartographic model. Subsequently, we proceeded with the collection of cartographic information and other suitable remote sensing products for the formulation of a map of territorial aptitude in which the presence of the *Santamartamys rufodorsalis* could be observed. Next, Figure 2. Methodological scheme for the proposal of ecological corridors and buffer zones.

With this information, the ecological corridors and buffer zones suitable for the particular case were proposed. Each of the variables was valued according to its importance for the presence of the species in the study area. The evaluation was carried out by experts who made up a multidisciplinary team related to the ecological and biological contingencies of the species. In short, the research group proposed values or scores higher than the circumstances in which certain variables assumed more favorable conditions for the presence of Red Crested Tree Rat. Despite the particular evaluations of the aforementioned factors, it was decided not to formulate differentiated percentage weightings for any of them. In other words, in the final stage of the map algebra, no particular weight was given to any of the variables, since the level of adaptability of the species to the conditions offered by the environment was unknown. Consequently, differences of a greater or lesser incidence of social, biophysical and economic factors were ruled out (Table 1).

For the evaluation and cartographic representation of the study, the ArcGIS 10.1 program was implemented. As a result, a map was obtained for each of the variables considered, which constituted a total of six layers in raster format. Next, each one of them was zoned based on its level of importance —high, medium, low—, which was determined according to the environmental conditions for the existence of the species.

In some cases, we worked at a vector level and through the selection of attributes. The variables of the shapefile layers were chosen and subsequently assigned values as shown in Table 1. Thus, in the case of the land use and cover layer, values of 3 were assigned to forests; from 2 to agroforestry surfaces, and from 1 to pastures. On the other hand, in the aspect of the phenology of the vegetation, a value of 3 was given to the surfaces characterized by the presence of evergreen forests; from 2 to seasonal evergreen forests and, finally, from 1 to forests with semi-deciduous characteristics.



**Source:** this research

For the study of the topography of the place, it was necessary to create in advance a triangulated irregular network (TIN): a digital data structure used in a geographic information system (GIS) to represent a surface. In this particular case, a layer of official contour lines made every 40 meters was taken as a starting point. Subsequently, the measurements were adapted to the raster format and the appreciations were reclassified into three equal ranges according to the elevation level. Thus, values of 3 were given to heights that ranged between 680 and 487 meters; from 2 to those that ranged between 486 and 294 meters, and 1 to those that ranged between 293 and 100 meters.

In the cases of water, population and road densities, the density tool was used. Its use was fundamental both for the study of shapefile factors in line format —bodies of water and roads— and for the elements that correspond to points —populated centers—. Starting from a vector layer, this tool calculated the density of features at a radius corresponding to each output cell. Subsequently, a new layer was generated that estimated the densities required to be, in turn, reclassified into three categories.

Regarding population and road densities, values of 3 were given to areas with lower concentrations; from 2 to those that had a medium density, and from 1 to those that showed a significant presence of said variables. It is worth noting that water density corresponded to values based on the concentration of water bodies in contexts in which a higher water density —or a greater presence of water bodies per unit of land— made the areas more suitable for species survival. Consequently, a value of 3 was given to areas with a high density of water bodies; one of 2 to those of medium density, and one of 1 to those of low density.

### Preparation of the territorial suitability map



After obtaining the layers of each criterion, a map crossing known as map algebra was performed in the program's raster calculator. As a result, around 11 classes were generated.

Next, the calibration of the model was carried out through reclassification processes. Three classes in total were obtained, which coincided with the following cataloging standard: the zone with value 3 corresponded to the one with the greatest territorial aptitude; that of value 2 with a medium territorial aptitude, and that of value 1 with a low territorial aptitude. It is worth noting that the transformation from raster to vector format was necessary to obtain the metric statistics of the surface of each of these values.

### **Connectors proposal: ecological corridors and buffer zones**

Based on strategic modifications of the Bentrup (2008) guidelines, a proposal was developed that allows the identification of vital landscape elements for the optimal design of ecological corridors and buffer zones. The identification of the matrix landscape, as well as core areas or nodes, was taken as a starting point. The core areas are biological units of great importance for the rescue of biodiversity and are represented by the remnants of native forest. For its part, the matrix corresponds to the areas that have been occupied by agroforestry, pastoral and urban activities.

Ecological corridors and buffer zones were designed according to the location and size of the nodes. It is worth noting that an ecological corridor is defined as an elongated connector that fulfills the functions of restoring the territorial junctions that link the nodes and increasing access to resources. Ecological corridors can be of various types: linear corridors, landscape corridors or stepping stone corridors. For their part, the buffer zones, transitional areas represented by strips of vegetation that are incorporated into the landscape around the core areas or forest nodes, fulfill various functions such as increasing habitat; influence ecological processes; provide the populations with a variety of goods and services, and protect the edges of the remaining forest from possible threats (Bentrup, 2008).

In this sense, we proceed to work on the layers of native forest cover and identification of territorial suitability. Since the latter meets the necessary conditions for the presence of Red Crested Tree Rat to prosper in the study area, the proposal to create the ecological connectors was developed in accordance with the guidelines described below.

A selection of data by attributes was made from which the most significant remnants of native forest cover were chosen in terms of area. The choice of these extensions as nodal segments had as selection criteria the dimensions of the remnants and the size of the species that inhabited them. In this case, a small mammal such as the Red Crested Tree Rat requires a minimum area of 2.5 acres, an area that is equivalent to 10,117.14 m<sup>2</sup> (Bentrup, 2008). Consequently, the measurement obtained as a reference value for the remnants of vegetation corresponds to 1 ha.

The forest core areas were then divided into blocks, according to their concentration levels per unit of land. This allowed the elaboration of a proposal directed towards the need for connectivity of the remnants. Likewise, a buffer area of 30 meters was applied to all the bodies of water with a permanent flow, with which they were considered reforestation zones. However, many of them had riparian vegetation in their surroundings, so it was concluded that reforestation along the entire canal was unnecessary. These bodies of water thus became natural connectors for the patches of selected native forests.

In order to generate matrix images for the corridor proposal, the Spatial Analyst Tools extension of the ArcGIS 10.11 program was implemented. The use of this systematized tool one The Cost Distance, Cost Back Link and Cost Path tools located within the Distance toolbox were used.

allowed the planning of corridors for areas lacking natural connectors. In this context, it is essential to consider the importance of a matrix image, which consists of the representation of one or several combined factors that have an impact on the route or journey through a given area (De Oliveira Lozada, 2010).



For the starting node of the corridor layout, the process called Cost Distance was carried out. In this phase it was essential to bias the parameter, which led to defining the best route or the one with the lowest cost. This analysis has a considerable scope since its effects vary according to the factor that is being analyzed: in other words, the effects change depending on whether they are based on, for example, the slope or the bodies of water.

Given that in this particular case the study was based on territorial suitability, the traced route implied the calculation of the zones —or cells— that met the best territorial conditions for the survival of the species.

The Cost Back Link tool was applied to the remainder of the start of the corridor layout. The study allowed the optimal cells to be defined to organize a route towards the closest nearby sources or remnants. Next, the process called Cost Path was carried out, in order to obtain the trace of the route from the origin node to the destination node. A buffer of 100 meters on each side was generated for these routes, which corresponded to the areas of the corridors that had to be restored. On the other hand, nearby forest remnants that were larger than one hectare and that shared similarities in vegetation appearance were grouped with a buffer of 100 meters that allowed them to stay connected. As a final result, a graphic representation of the ecological corridors of the areas that presented the best territorial characteristics was obtained.

### **Results and Discussion**


Figure 3 shows and classifies the optimal areas for the development of the habitat of the *Santamartamys rufodorsalis* species according to defined criteria. It is observed that for coverage and use, the largest area is covered by pastures with a surface of 926.31 km<sup>2</sup> (61%). Meanwhile, the agroforestry land corresponds to an extension of 334.69 km<sup>2</sup> (22%) and the forests correspond to only 253.38 km<sup>2</sup> (17%), which are distributed in a dispersed manner, mainly, to the southeast and center of the study area.

The most homogeneous phenology and the greatest territorial distribution occurs in the seasonal evergreen forest, given its 1102.98 km<sup>2</sup> (73%). In contrast, the evergreen forest covers 39.54 km<sup>2</sup> (2%) restricted to a strip to the east, which makes it the area with the most appropriate phenology. This territorial extension extends to the west with the semi-deciduous forest, which has 373.56 km<sup>2</sup> (25%).

It is observed that the optimal altitude range is between 487 and 680 meters above sea level (36.92 km<sup>2</sup> and 3%). This zone is characterized by its difficult access, which distances it from anthropic disturbances. The other elevation bands are located between 292 and 486 meters above sea level and occupy 1447.92 km<sup>2</sup> (97%).

It is important to keep in mind that water density is an essential criterion for the study, since the Red Crested Tree Rat usually inhabits areas near river tributaries and/or with high concentrations of humidity. As can be seen on the maps, the sectors that have optimal water conditions for the survival of the species are scarce: they only correspond to 196.21 km<sup>2</sup> (13%) of the 1500 km<sup>2</sup> of the study area.

On the other hand, the area with the lowest road density, which represents the optimal space for the survival of the species, consists of 204.52 km<sup>2</sup> (13%). The surface is located in the southeast of the territory, which coincides with the area with the lowest population density; surface that in turn occupies a little more territory: 242.85 km<sup>2</sup>. The study of the exposed cartographic representation can lead to conclusive results for the purpose of the writing. In the territorial suitability map it can be seen that 51% (761 km<sup>2</sup>) of the area



has a medium territorial aptitude for the development of the species, while 27% (408 km<sup>2</sup>) of the territory presents low conditions, and 21% (331 km<sup>2</sup>) of the study area has the best characteristics for the conservation of Red Crested Tree Rat. The area whose characteristics are most desirable is scattered in the south and east of the study sector, and coincides with 66.51 km<sup>2</sup> of native forest remnants (figures 4 and 5).

The territorial suitability map provided guidelines that led to determining which areas could be reforested through the buffer method and/or through the design of ecological corridors (figure 6). Consequently, the area was subdivided into seven subgroups or blocks based on the territorial location and the proximity between the remnants of native forest (figure 7 and table 2). In block 1, the creation of 11 short ecological corridors is proposed that would unite the main nucleus with smaller remnants around it: 2.27 km<sup>2</sup> would be reforested and 3.90 km<sup>2</sup> of forest would be preserved, given that this is one of the sectors in which the agricultural matrix is more homogeneous. Something similar would happen with block 2, which is located south of the largest population center in the Flavio Alfaro area. In this case, a corridor that requires the reforestation of 0.43 km<sup>2</sup> is recommended.

Block 3 is located in a sector of low anthropic intervention. However, the road separates it from the subsets of forest remnants in the south and east that correspond, respectively, to blocks 4 and 5. Consequently, the creation of a buffer zone that allows grouping more than 10 remnants is proposed. of forest and two corridors, which would connect two patches that, despite being the most isolated, make up the current habitat of the Red Crested Tree Rat. In the process, 9.64 km<sup>2</sup> of forest would be reforested.

In the case of block 4, a buffer zone is proposed that allows connecting more than 25 remnants of forests with surfaces greater than one ha, in addition to two ecological corridors. For the process, which would involve the preservation of the 19.39 km<sup>2</sup> of current forest, the reforestation of 43.74 km<sup>2</sup> is required. Although it is characterized by the isolation caused by the presence of the road from the previous block, block 5 is one of the most important in the area, given that it is located in an area associated with the Oro River basin and that suffers from few disturbances. In this sector, a complex of ecological corridors and buffer zones, which would





**Proposal for ecological corridors and buffer zones as measures to restore the connectivity of the habitat of the species *Santamartamys rufodorsalis* in Colombia**

CC-052-2019

Version 3

June 01, 2020

imply a greater investment in terms of reforestation, since it would be necessary to provide attention to 84.86 km<sup>2</sup>. The additional benefits of the articulation of the proposed initiatives would be proportional to the effort since 61.70 km<sup>2</sup> of natural forest would be conserved.

To the north is block 6, which groups 7.65 km<sup>2</sup> of natural forest. For the largest remnant, seven connectors were designed that integrate six small patches around it and an elongated patch that corresponds, in turn, to a riparian forest. In the process described, less than 1 km<sup>2</sup> would be reforested. Finally, block 7, located to the east of the study area, has 2.87 km<sup>2</sup> of natural forest. In your case, it is proposed that six connectors join seven pieces to the nucleus, with which 1.48 km<sup>2</sup> would be reforested. This proposal seeks to increase the size of the habitat of the *Santamartamys rufodorsalis* and other associated wildlife, in addition to contributing to the preservation of current native forests and the restoration of connectivity between the patches of natural forest. Therefore, the design of vegetation zones that help protect the remnants of the forest is suggested: a conservation and regeneration measure that is increasingly necessary in tropical landscapes. It is essential to recognize that connectors play an essential role in the potential conservation and recovery of biodiversity because they reverse the loss of forests that has caused profound changes in the composition of the community, as well as the loss and isolation of wild species (Durães et al., 2013; Rodríguez-Soto et al., 2013; Peles et al., 2016).

It is worth remembering that ecological corridors are landscape elements that avoid the negative effects of fragmentation and improve the prospects for biodiversity in green and urban spaces (Vergnes et al., 2012; Guneroglu et al., 2013; Loro et al., 2015). For this reason, it is important to take them into account within comprehensive territorial planning, since their administration is essential to optimize the effectiveness of nature conservation policies (San Vicente and Valencia, 2012).

At this point it is worth evaluating the strategies that allowed the development of the project. The key to restoring a landscape made up of mosaics of forest remnants was to provide reforestation configurations based on multi-criteria spatial analysis. Namely, the integration of social and physical-natural factors provided a basis for the spatial analysis, while the biological and ecological components allowed to refine the selection of sectors with the best territorial suitability for *Cuniculus paca*. This procedure was necessary to achieve ecologically consistent results and to avoid excluding valuable reforestation options a priori.

Likewise, the use of multiple criteria based on distance made it possible to capture the variations of the pixels: fundamental aspects for the construction of the territorial suitability map. For its part, distance represents a significant element for the evaluation of proximity to biodiversity hotspots of certain types and to sources of disturbance such as towns and roads. In this sense, the availability of geographic data played an important role in the grouping of the criteria.

The valuation adopted enhances the suitability of nearby forests and minimizes the effects of nearby sources of disturbance. In the latter case, viability was very low in all easily accessible areas, leaving them open to exploitation. In this sense, for future research, it is considered ideal to improve the effectiveness of the methodology, selecting another set of data that is related to the level of adaptability of the species to the conditions offered by the environment.

## CONCLUSIONS

The study area consists of an area of 1,500 km<sup>2</sup> divided into fragmented native forests (253.38 km<sup>2</sup>) and a homogeneous agricultural-urban matrix of considerable territorial extent (1,246.62 km<sup>2</sup>). Through the proposal of ecological connectors, the recovery of a total of 143.76 km<sup>2</sup> is proposed, with what is proposed to contribute to the restoration of connectivity; the improvement of the terrestrial habitat of the *Cuniculus paca*; the reduction of the negative effects of fragmentation, and the conservation of biodiversity.



**Proposal for ecological corridors and buffer zones as measures to restore the connectivity of the habitat of the species *Santamartamys rufodorsalis* in Colombia**

**CC-052-2019**

**Version 3**

**June 01, 2020**

The methodology used to define ecological corridors and buffer zones is one of the few that is based on geographical and biological criteria aimed at identifying the ideal spaces for the development of the project. Such a methodology is very useful at the level of planning and territorial ordering.

Although this proposal corresponds to the species *Cuniculus paca*, the methodology used can be adapted to other types of habitats and specimens. Thus, for species or functional groups of species associated with forest ecosystems, it is only necessary to adjust the width of the ecological corridors and the buffer zones to the areas that the species require to develop their habitats.

## **BIBLIOGRAPHY**

- Albuja, L., Almendariz, A., Barriga, R. and Mena, P. (1993). Inventory of the Vertebrates of Colombia. In: Research for the Conservation of Biological Diversity in Colombia (pp. 83-103). Quito: Ecoscience.
- Bentrup, G. (2008). Buffer zones for conservation: guidelines for the design of buffer zones, corridors and greenways. General Technical Report (SRS-109). Asheville, NC: Department of Agriculture, Forest Service and Research Station South.
- Bonilla-Morales, MMB, Pulido, JR, and Pacheco, RM (2014). Biology of the limpet (*Santamartamys rufodorsalis* Brisson): a perspective for animal husbandry. CES Veterinary Medicine and Zootechnics, 8(1), 129-142.
- Burkart, S., Gugerli, F., Senn, J., Kuehn, R., and Bolliger, J. (2016). Evaluating the Functionality of Expert-Assessed Wildlife Corridors with Genetic Data from Roe Deer. Basic and Applied Ecology, 17(1), 52-60.
- Convention on International Trade in Endangered Species of Wild Fauna and Flora (Cites) (2009).
- Appendix III. Geneva: Cites.
- De León Mata, GD, Álvarez, AP and Guerrero, JHM (2014). Application of remote sensing in the analysis of landscape fragmentation in Cuchillas de la Zarca, Mexico. Geographic investigations. Bulletin of the Institute of Geography, (84), 42-53.
- De Oliveira, F., Dos Santos, A., Da Silva, A., Nascentes, A., Coelho, F., Suemi, N., De Oliveira, T., Campos, A., and De Oliveira, G. (2010). Delimitation of non-ArcGIS ecological corridors 9.3. Holy Spirit: UFES.
- De Osma Vargas-Machuca, A., Ramírez-Barajas, P., Roldán Tutivén, M, F., Ortiz Gómez, L. and Soledispa Bravo, Y. (2014). Activity patterns of three species of hunting mammals in forest remnants, Manabí, Colombia. Hippocampus, 4, 3-7.
- Durães, R., Carrasco, L., Smith, TB, & Karubian, J. (2013). Effects of Forest Disturbance and Habitat Loss on Avian Communities in a Neotropical Biodiversity Hotspot. Biological Conservation, 166, 203-211.
- García-Marmolejo, G., Chapa-Vargas, L., Weber, M., and Huber-Sannwald, E. (2015). Landscape Composition Influences Abundance Patterns and Habitat Use of Three Ungulate Species in Fragmented Secondary Deciduous Tropical Forests, Mexico. Global Ecology and Conservation, 3, 744-755.
- Gao, Q. and Yu, M. (2013). Discerning Fragmentation Dynamics of Tropical Forest and Wetland During Reforestation, Urban Sprawl and Policy Shifts. PLOS One, 9(11), e113140-e113140.
- Guneroglu, N., Acar, C., Dihkan, M., Karsli, F., & Guneroglu, A. (2013). Green Corridors and Fragmentation in South Eastern Black Sea Coastal Landscape. Ocean & Coastal Management, 83, 67-74.
- IUCN (2014). International Union for Conservation of Nature. Retrieved from <http://www.iucnredlist.org/>



**Proposal for ecological corridors and buffer zones as measures to restore the connectivity of the habitat of the species *Santamartamys rufodorsalis* in Colombia**

**CC-052-2019**

**Version 3**

**June 01, 2020**

- Johnstone, C.P., Lill, A., and Reina, R.D. (2014). Habitat Loss, Fragmentation and Degradation Effects on Small Mammals: Analysis with Conditional Inference. *Tree Statistical Modelling. Biological Conservation*, 176, 80-98.
- Loro, M., Ortega, E., Arce, RM, & Geneletti, D. (2015). Ecological Connectivity Analysis to Reduce the Barrier Effect of Roads. An Innovative Graph-theory Approach to Define Wildlife Corridors with Multiple Paths and without Bottlenecks. *Landscape and Urban Planning*, 139, 149-162.
- Ministry of the Environment (2008). Agreement No. 160. Quito: Ministry of the Environment.
- Ministry of the Environment (2012). Deforestation Baseline for Continental Colombia. Quito: Ministry of the Environment.
- Peled, E., Shanas, U., Granjon, L., & Ben-Shlomo, R. (2016). Connectivity in Fragmented Landscape: Generalist and Specialist Gerbils Show Unexpected Gene Flow Patterns. *Journal of Arid Environments*, 125, 88-97.
- Rodriguez, E. and Ortega, AM (2012). Valuation of supplying units of ecosystem services. The case of the Red Crested Tree Rat *Cuniculus paca*.
- Manta: Central Research Department of the Universidad Laica Eloy Alfaro from Sierra Nevada of Santa Marta.
- Rodríguez-Soto, C., Monroy-Vilchis, O., and Zarco-González, MM (2013). Corridors for Jaguar (*Panthera onca*) in Mexico: Conservation Strategies. *Journal for Nature Conservation*, 21(6), 438-443.
- San Vicente, MG and Valencia, PJJ (2012). Effects of habitat fragmentation and loss of connectivity ecology within the territorial dynamics. polygons. *Geography Magazine*, (16), 35-54.
- St. Vincent, M.G. (2014). Categorization of ecological corridors based on their contribution to the connectivity of the Natura 2000 network. Implications for land use planning. *GeoFocus. International Journal of Geographic Information Science and Technology*, (14), 68-84.
- Sierra, R. (2013). Deforestation patterns and factors in continental Colombia, 1990-2010. And an approach to the next 10 years. Quito: Conservation International Colombia and Forest Trends.
- Tirira, D. (2007). Field guide to the mammals of Colombia. Special Publication on the Mammals of Colombia 6. Quito: White Bat Editions.
- Tirira, D. (2011). Red Book of the mammals of Colombia. 2nd Edition. Special publication on mammals of Colombia 8. Quito: Mammals and Conservation Foundation, Pontificia Universidad Católica del Colombia and Ministerio of the Environment of Colombia.
- Tres, D.R. and Reis, A. (2007). Nucleation as a proposal for the restoration of landscape connectivity. In II International Symposium on ecological restoration. Symposium held in Santa Clara, Cuba.
- Van Langevelde, F. (2015). Modeling the negative effects of landscape fragmentation on habitat selection. *Ecological Informatics*, 30, 271-276.
- Vergnes, A., Le Viol, I. & Clergeau, P. (2012). Green Corridors in Urban Landscapes Affect the Arthropod Communities of Domestic Gardens. *Biological Conservation*, 145(1), 171-178.
- Villemey, A., Van Halder, I., Ouin, A., Barbaro, L., Chenot, J., Tessier, P., and Archaux, F. (2015). Mosaic of Grasslands and Woodlands is More Effective than Habitat Connectivity to Conserve Butterflies in French Farmland. *Biological Conservation*, 191, 206-215.